


RETTOANGOLO



p = perimetro b

$$p = 2(b + h)$$

$$b = \frac{p - 2h}{2}$$

$$h = \frac{p - 2b}{2}$$

$$A = b \times h$$

$$b = \frac{A}{h}$$

$$h = \frac{A}{b}$$



p = perimetro b

$$p = 2(b + c)$$

$$b = \frac{p - 2c}{2}$$


$$c = \frac{p - 2b}{2}$$

$$A = b \times h$$

$$b = \frac{A}{h}$$

$$h = \frac{A}{b}$$

QUADRATO



p = perimetro ℓ

$$p = 4\ell$$


$$\ell = p : 4$$

$$A = \ell^2$$

$$\ell = \sqrt{A}$$

$$A = \frac{d^2}{2} \quad d = \sqrt{A \times 2}$$


ROMBO



$$p = 4\ell \quad \ell = p : 4$$

$$A = \frac{d_1 \times d_2}{2} \quad d_1 = \frac{2 \times A}{d_2}$$


TRAPEZIO



$$A = \frac{(b_1 + b_2) \times h}{2}$$

$$h = \frac{A \times 2}{b_1 + b_2} \quad b_1 + b_2 = \frac{A \times 2}{h}$$


TRIANGolo



$$A = \frac{b \times h}{2}$$

$$h = \frac{A \times 2}{b} \quad b = \frac{A \times 2}{h}$$

TRIANGolo RETTANGOLO



$$A = \frac{c_1 \times c_2}{2} \quad c_1 = \frac{A \times 2}{c_2}$$

$$h_i = \frac{c_1 \times c_2}{i}$$